

# Hidden structures of holographic correlators

Connor Behan

Oxford Mathematical Institute

2021-08-12

[2101.04114] with P. Ferrero, X. Zhou

[2103.15830] with L. F. Alday, P. Ferrero, X. Zhou

# Types of hidden symmetries

	Chiral	Hidden conformal	Parisi-Sourlas
3d $\mathcal{N} = 8$ ABJM			
4d $\mathcal{N} = 4$ SYM			
6d $\mathcal{N} = (2, 0)$			
3d $\mathcal{N} = 3$ flavoured ABJM			
4d $\mathcal{N} = 2$ Argyres-Douglas			
5d $\mathcal{N} = 1$ Seiberg			
6d $\mathcal{N} = (1, 0)$ E-string			

# Types of hidden symmetries

	Chiral	Hidden conformal	Parisi-Sourlas
3d $\mathcal{N} = 8$ ABJM	Red		
4d $\mathcal{N} = 4$ SYM	Green		
6d $\mathcal{N} = (2, 0)$	Green		
3d $\mathcal{N} = 3$ flavoured ABJM	Red		
4d $\mathcal{N} = 2$ Argyres-Douglas	Green		
5d $\mathcal{N} = 1$ Seiberg	Red		
6d $\mathcal{N} = (1, 0)$ E-string	Red		

# Types of hidden symmetries

	Chiral	Hidden conformal	Parisi-Sourlas
3d $\mathcal{N} = 8$ ABJM	Red	Red	
4d $\mathcal{N} = 4$ SYM	Green	Green	
6d $\mathcal{N} = (2, 0)$	Green	Red	
3d $\mathcal{N} = 3$ flavoured ABJM	Red	Red	
4d $\mathcal{N} = 2$ Argyres-Douglas	Green	Green	
5d $\mathcal{N} = 1$ Seiberg	Red	Red	
6d $\mathcal{N} = (1, 0)$ E-string	Red	Red	

# Types of hidden symmetries

	Chiral	Hidden conformal	Parisi-Sourlas
3d $\mathcal{N} = 8$ ABJM	Red	Red	Green
4d $\mathcal{N} = 4$ SYM	Green	Green	Green
6d $\mathcal{N} = (2, 0)$	Green	Red	Green
3d $\mathcal{N} = 3$ flavoured ABJM	Red	Red	Green
4d $\mathcal{N} = 2$ Argyres-Douglas	Green	Green	Green
5d $\mathcal{N} = 1$ Seiberg	Red	Red	Green
6d $\mathcal{N} = (1, 0)$ E-string	Red	Red	Green

# Types of hidden symmetries

	Chiral	Hidden conformal	Parisi-Sourlas
3d $\mathcal{N} = 8$ ABJM	Yellow	Red	Green
4d $\mathcal{N} = 4$ SYM	Green	Green	Green
6d $\mathcal{N} = (2, 0)$	Green	Red	Green
3d $\mathcal{N} = 3$ flavoured ABJM	Red	Red	Green
4d $\mathcal{N} = 2$ Argyres-Douglas	Green	Green	Green
5d $\mathcal{N} = 1$ Seiberg	Red	Red	Green
6d $\mathcal{N} = (1, 0)$ E-string	Red	Red	Green

# Types of hidden symmetries

	Chiral	Hidden conformal	Parisi-Sourlas
3d $\mathcal{N} = 8$ ABJM	Yellow	Red	Green
4d $\mathcal{N} = 4$ SYM	Green	Green	Green
6d $\mathcal{N} = (2, 0)$	Green	Red	Green
3d $\mathcal{N} = 3$ flavoured ABJM	Yellow	Red	Green
4d $\mathcal{N} = 2$ Argyres-Douglas	Green	Green	Green
5d $\mathcal{N} = 1$ Seiberg	Red	Red	Green
6d $\mathcal{N} = (1, 0)$ E-string	Red	Red	Green

Consider a theory with  $SL(2)$  conformal,  $SU(2)$  global symmetry:

$$\mathcal{O}(z, v) = \mathcal{O}^{\alpha_1 \dots \alpha_{2j}}(z) v_{\alpha_1} \dots v_{\alpha_{2j}}.$$

4pt functions depend on cross ratios:

$$\chi = \frac{z_{12} z_{34}}{z_{13} z_{24}}, \quad \alpha = \frac{v_{13} v_{24}}{v_{12} v_{34}}.$$



Consider a theory with  $SL(2)$  conformal,  $SU(2)$  global symmetry:

$$\mathcal{O}(z, v) = \mathcal{O}^{\alpha_1 \dots \alpha_{2j}}(z) v_{\alpha_1} \dots v_{\alpha_{2j}}.$$

4pt functions depend on cross ratios:

$$\chi = \frac{z_{12} z_{34}}{z_{13} z_{24}}, \quad \alpha = \frac{v_{13} v_{24}}{v_{12} v_{34}}.$$

One common convention is:

$$\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle = \left( \frac{z_{24}}{z_{14}} \right)^{h_{12}} \left( \frac{z_{14}}{z_{13}} \right)^{h_{34}} \frac{1}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \mathcal{G}(\chi).$$

Consider a theory with  $SL(2)$  conformal,  $SU(2)$  global symmetry:

$$\mathcal{O}(z, v) = \mathcal{O}^{\alpha_1 \dots \alpha_{2j}}(z) v_{\alpha_1} \dots v_{\alpha_{2j}}.$$

4pt functions depend on cross ratios:

$$\chi = \frac{z_{12} z_{34}}{z_{13} z_{24}}, \quad \alpha = \frac{v_{13} v_{24}}{v_{12} v_{34}}.$$

One common convention is:

$$\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle = \left( \frac{z_{24}}{z_{14}} \right)^{h_{12}} \left( \frac{z_{14}}{z_{13}} \right)^{h_{34}} \left( \frac{v_{14}}{v_{24}} \right)^{j_{12}} \left( \frac{v_{13}}{v_{14}} \right)^{j_{34}} \frac{v_{12}^{j_1+j_2} v_{34}^{j_3+j_4}}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \mathcal{G}(\chi; \alpha).$$

Consider a theory with  $SL(2)$  conformal,  $SU(2)$  global symmetry:

$$\mathcal{O}(z, v) = \mathcal{O}^{\alpha_1 \dots \alpha_{2j}}(z) v_{\alpha_1} \dots v_{\alpha_{2j}}.$$

4pt functions depend on cross ratios:

$$\chi = \frac{z_{12} z_{34}}{z_{13} z_{24}}, \quad \alpha = \frac{v_{13} v_{24}}{v_{12} v_{34}}.$$

One common convention is:

$$\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle = \left( \frac{z_{24}}{z_{14}} \right)^{h_{12}} \left( \frac{z_{14}}{z_{13}} \right)^{h_{34}} \left( \frac{v_{14}}{v_{24}} \right)^{j_{12}} \left( \frac{v_{13}}{v_{14}} \right)^{j_{34}} \frac{v_{12}^{j_1+j_2} v_{34}^{j_3+j_4}}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \mathcal{G}(\chi; \alpha).$$

Another uses **extremality**  $\mathcal{E} = 2\min(j_i)$  or  $\sum_i j_i - 2\max(j_i)$ .

$$\sqrt{\frac{v_{34}^{j_3+j_4-j_1-j_2}}{34}} \sqrt{\frac{v_{24}^{j_2+j_4-j_1-j_3}}{24}} \sqrt{\frac{v_{23}^{j_1+j_2+j_3-j_4-\mathcal{E}}}{23}} v_{14}^{2j_1-\mathcal{E}} (v_{12} v_{34})^{\mathcal{E}}.$$

Consider a theory with  $SL(2)$  conformal,  $SU(2)$  global symmetry:

$$\mathcal{O}(z, v) = \mathcal{O}^{\alpha_1 \dots \alpha_{2j}}(z) v_{\alpha_1} \dots v_{\alpha_{2j}}.$$

4pt functions depend on cross ratios:

$$\chi = \frac{z_{12} z_{34}}{z_{13} z_{24}}, \quad \alpha = \frac{v_{13} v_{24}}{v_{12} v_{34}}.$$

One common convention is:

$$\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle = \left( \frac{z_{24}}{z_{14}} \right)^{h_{12}} \left( \frac{z_{14}}{z_{13}} \right)^{h_{34}} \left( \frac{v_{14}}{v_{24}} \right)^{j_{12}} \left( \frac{v_{13}}{v_{14}} \right)^{j_{34}} \frac{v_{12}^{j_1+j_2} v_{34}^{j_3+j_4}}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \mathcal{G}(\chi; \alpha).$$

Another uses **extremality**  $\mathcal{E} = 2\min(j_i)$  or  $\sum_i j_i - 2\max(j_i)$ .

$$\left[ \frac{v_{34}}{x_{34}^{2\epsilon}} \right]^{j_3+j_4-j_1-j_2} \left[ \frac{v_{24}}{x_{24}^{2\epsilon}} \right]^{j_2+j_4-j_1-j_3} \left[ \frac{v_{23}}{x_{23}^{2\epsilon}} \right]^{j_1+j_2+j_3-j_4-\mathcal{E}} \left[ \frac{v_{14}}{x_{14}^{2\epsilon}} \right]^{2j_1-\mathcal{E}} \left[ \frac{v_{12} v_{34}}{x_{12}^{2\epsilon} x_{34}^{2\epsilon}} \right]^{\mathcal{E}}$$

Consider a theory with  $SL(2)$  conformal,  $SU(2)$  global symmetry:

$$\mathcal{O}(z, v) = \mathcal{O}^{\alpha_1 \dots \alpha_{2j}}(z) v_{\alpha_1} \dots v_{\alpha_{2j}}.$$

4pt functions depend on cross ratios:

$$\chi = \frac{z_{12} z_{34}}{z_{13} z_{24}}, \quad \alpha = \frac{v_{13} v_{24}}{v_{12} v_{34}}.$$

One common convention is:

$$\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle = \left( \frac{z_{24}}{z_{14}} \right)^{h_{12}} \left( \frac{z_{14}}{z_{13}} \right)^{h_{34}} \left( \frac{v_{14}}{v_{24}} \right)^{j_{12}} \left( \frac{v_{13}}{v_{14}} \right)^{j_{34}} \frac{v_{12}^{j_1+j_2} v_{34}^{j_3+j_4}}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \mathcal{G}(\chi; \alpha).$$

Another uses **extremality**  $\mathcal{E} = 2\min(j_i)$  or  $\sum_i j_i - 2\max(j_i)$ .

$$\left[ \frac{v_{34}}{x_{34}^{2\epsilon}} \right]^{j_3+j_4-j_1-j_2} \left[ \frac{v_{24}}{x_{24}^{2\epsilon}} \right]^{j_2+j_4-j_1-j_3} \left[ \frac{v_{23}}{x_{23}^{2\epsilon}} \right]^{j_1+j_2+j_3-j_4-\mathcal{E}} \left[ \frac{v_{14}}{x_{14}^{2\epsilon}} \right]^{2j_1-\mathcal{E}} \left[ \frac{v_{12} v_{34}}{x_{12}^{2\epsilon} x_{34}^{2\epsilon}} \right]^{\mathcal{E}}$$

For  $d > 2$ , use  $U = \chi\chi'$  and  $V = (1 - \chi)(1 - \chi')$ .

# Maximal and half-maximal SUSY

With 16 Qs, R symmetry is  $SO(5)$ ,  $SO(6)$  or  $SO(8)$ .

$$\mathcal{O}(x, t) = \mathcal{O}^{l_1 \dots l_k}(x) t_{l_1} \dots t_{l_k}, \quad \Delta = \epsilon k, \quad \epsilon = \frac{d-2}{2}$$
$$\sigma = \frac{t_{13} t_{24}}{t_{12} t_{34}} = \alpha \alpha', \quad \tau = \frac{t_{14} t_{23}}{t_{12} t_{34}} = (1 - \alpha)(1 - \alpha')$$

# Maximal and half-maximal SUSY

With **16 Qs**, R symmetry is  $SO(5)$ ,  $SO(6)$  or  $SO(8)$ .

$$\mathcal{O}(x, t) = \mathcal{O}^{l_1 \dots l_k}(x) t_{l_1} \dots t_{l_k}, \quad \Delta = \epsilon k, \quad \epsilon = \frac{d-2}{2}$$
$$\sigma = \frac{t_{13} t_{24}}{t_{12} t_{34}} = \alpha \alpha', \quad \tau = \frac{t_{14} t_{23}}{t_{12} t_{34}} = (1 - \alpha)(1 - \alpha')$$

One can use a brane to break this to **8 Qs** and  $SU(2)_L \times SU(2)_R$ .

$$\mathcal{O}(x, v, \bar{v}) = \mathcal{O}_{\beta_1 \dots \beta_{k-2}}^{\alpha_1 \dots \alpha_k}(x) v_{\alpha_1} \dots v_{\alpha_k} \bar{v}^{\beta_1} \dots \bar{v}^{\beta_{k-2}}, \quad \Delta = \epsilon k$$
$$\alpha = \frac{v_{12} v_{34}}{v_{13} v_{24}}, \quad \beta = \frac{\bar{v}_{12} \bar{v}_{34}}{\bar{v}_{13} \bar{v}_{24}}, \quad \sigma = \alpha \beta, \quad \tau = (1 - \alpha)(1 - \beta)$$

# Maximal and half-maximal SUSY

With **16 Qs**, R symmetry is  $SO(5)$ ,  $SO(6)$  or  $SO(8)$ .

$$\mathcal{O}(x, t) = \mathcal{O}^{l_1 \dots l_k}(x) t_{l_1} \dots t_{l_k}, \quad \Delta = \epsilon k, \quad \epsilon = \frac{d-2}{2}$$
$$\sigma = \frac{t_{13} t_{24}}{t_{12} t_{34}} = \alpha \alpha', \quad \tau = \frac{t_{14} t_{23}}{t_{12} t_{34}} = (1 - \alpha)(1 - \alpha')$$

One can use a brane to break this to **8 Qs** and  $SU(2)_L \times SU(2)_R$ .

$$\mathcal{O}(x, v, \bar{v}) = \mathcal{O}_{\beta_1 \dots \beta_{k-2}}^{\alpha_1 \dots \alpha_k}(x) v_{\alpha_1} \dots v_{\alpha_k} \bar{v}^{\beta_1} \dots \bar{v}^{\beta_{k-2}}, \quad \Delta = \epsilon k$$
$$\alpha = \frac{v_{12} v_{34}}{v_{13} v_{34}}, \quad \beta = \frac{\bar{v}_{12} \bar{v}_{34}}{\bar{v}_{13} \bar{v}_{34}}, \quad \sigma = \alpha \beta, \quad \tau = (1 - \alpha)(1 - \beta)$$

Four (two) identities for 16 Qs (8 Qs) [Dolan, Gallot, Sokatchev; [hep-th/0405180](http://hep-th/0405180)].

$$(\chi \partial_\chi - \epsilon \alpha \partial_\alpha) \mathcal{G} \Big|_{\alpha=\chi^{-1}} = 0$$



# Holography in Mellin space

Use Mandelstam variables instead of cross ratios [\[Mack; 0907.2407\]](#) .

$$\mathcal{G}(U, V) = \int_{-i\infty}^{i\infty} \frac{ds dt}{(4\pi i)^2} \frac{V^{\frac{\epsilon}{2}(k_1 - k_4 - \mathcal{E}) + \frac{t}{2}}}{U^{\frac{\epsilon}{2}(k_1 + k_2 - \mathcal{E}) - \frac{s}{2}}} \mathcal{M}(s, t) \Gamma\left[\frac{\Delta_1 + \Delta_2 - s}{2}\right] \Gamma\left[\frac{\Delta_3 + \Delta_4 - s}{2}\right] \\ \Gamma\left[\frac{\Delta_1 + \Delta_4 - t}{2}\right] \Gamma\left[\frac{\Delta_2 + \Delta_3 - t}{2}\right] \Gamma\left[\frac{\Delta_1 + \Delta_3 - u}{2}\right] \Gamma\left[\frac{\Delta_2 + \Delta_4 - u}{2}\right]$$

# Holography in Mellin space

Use Mandelstam variables instead of cross ratios [\[Mack; 0907.2407\]](#).

$$\mathcal{G}(U, V) = \int_{-i\infty}^{i\infty} \frac{dsdt}{(4\pi i)^2} \frac{V^{\frac{\epsilon}{2}(k_1 - k_4 - \mathcal{E}) + \frac{t}{2}}}{U^{\frac{\epsilon}{2}(k_1 + k_2 - \mathcal{E}) - \frac{s}{2}}} \mathcal{M}(s, t) \Gamma\left[\frac{\Delta_1 + \Delta_2 - s}{2}\right] \Gamma\left[\frac{\Delta_3 + \Delta_4 - s}{2}\right] \\ \Gamma\left[\frac{\Delta_1 + \Delta_4 - t}{2}\right] \Gamma\left[\frac{\Delta_2 + \Delta_3 - t}{2}\right] \Gamma\left[\frac{\Delta_1 + \Delta_3 - u}{2}\right] \Gamma\left[\frac{\Delta_2 + \Delta_4 - u}{2}\right]$$

Conformal blocks and Witten diagrams (for  $\tau = \Delta - \ell$ ) both become

$$\mathcal{M}_{\tau, \ell}(s, t) = \sum_{m=0}^{\infty} \frac{Q_{\ell, m}^{\tau}(t)}{m! \Gamma\left[\frac{\Delta_1 + \Delta_2 - \tau}{2} - m\right] \Gamma\left[\frac{\Delta_3 + \Delta_4 - \tau}{2} - m\right] (s - \tau - 2m)}.$$

# Holography in Mellin space

Use Mandelstam variables instead of cross ratios [Mack; 0907.2407].

$$\mathcal{G}(U, V) = \int_{-i\infty}^{i\infty} \frac{dsdt}{(4\pi i)^2} \frac{V^{\frac{\epsilon}{2}(k_1 - k_4 - \mathcal{E}) + \frac{t}{2}}}{U^{\frac{\epsilon}{2}(k_1 + k_2 - \mathcal{E}) - \frac{s}{2}}} \mathcal{M}(s, t) \Gamma \left[ \frac{\Delta_1 + \Delta_2 - s}{2} \right] \Gamma \left[ \frac{\Delta_3 + \Delta_4 - s}{2} \right] \\ \Gamma \left[ \frac{\Delta_1 + \Delta_4 - t}{2} \right] \Gamma \left[ \frac{\Delta_2 + \Delta_3 - t}{2} \right] \Gamma \left[ \frac{\Delta_1 + \Delta_3 - u}{2} \right] \Gamma \left[ \frac{\Delta_2 + \Delta_4 - u}{2} \right]$$

Conformal blocks and Witten diagrams (for  $\tau = \Delta - \ell$ ) both become

$$\mathcal{M}_{\tau, \ell}(s, t) = \sum_{m=0}^{\infty} \frac{Q_{\ell, m}^{\tau}(t)}{m! \Gamma \left[ \frac{\Delta_1 + \Delta_2 - \tau}{2} - m \right] \Gamma \left[ \frac{\Delta_3 + \Delta_4 - \tau}{2} - m \right] (s - \tau - 2m)}.$$

Add **contact terms** to single trace blocks in all channels.

$$\mathcal{M}(s, t) = \sum_{\mathcal{O}} C_{12\mathcal{O}} C_{34\mathcal{O}} \mathcal{M}_{\mathcal{O}}(s, t) + C_{14\mathcal{O}} C_{23\mathcal{O}} \mathcal{M}_{\mathcal{O}}(t, s) + C_{13\mathcal{O}} C_{24\mathcal{O}} \mathcal{M}_{\mathcal{O}}(u, t) \\ + P_{\ell_{\max} - 1}(s, t)$$

# Operator content

Instead of  $\mathcal{M}_{\mathcal{O}}(s, t)$ , use  $\mathcal{S}_{\mathcal{O}}(s, t; \alpha)$  which is a linear combination of  $\mathcal{Y}_j(\alpha)\mathcal{M}_{\tau, \ell}(s, t)$  for all components of the superconformal block.

# Operator content

Instead of  $\mathcal{M}_{\mathcal{O}}(s, t)$ , use  $\mathcal{S}_{\mathcal{O}}(s, t; \alpha)$  which is a linear combination of  $\mathcal{Y}_j(\alpha)\mathcal{M}_{\tau, \ell}(s, t)$  for all components of the superconformal block.

	$OSp(8 4)$	$PSU(2, 2 2)$	$OSp(8^* 4)$
$\ell = 0$	$B[0]_{p/2}^{[0,0,p,0]}$	$B\bar{B}[0, 0]_p^{[0,p,0]}$	$D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 2$	$Q^4 B[0]_{p/2}^{[0,0,p,0]}$	$Q^2 \bar{Q}^2 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^4 D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 0$	$Q^8 B[0]_{p/2}^{[0,0,p,0]}$	$Q^4 \bar{Q}^4 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^8 D[0, 0, 0]_{2p}^{[p,0]}$

# Operator content

Instead of  $\mathcal{M}_{\mathcal{O}}(s, t)$ , use  $\mathcal{S}_{\mathcal{O}}(s, t; \alpha)$  which is a linear combination of  $\mathcal{Y}_j(\alpha)\mathcal{M}_{\tau, \ell}(s, t)$  for all components of the superconformal block.

	$OSp(8 4)$	$PSU(2, 2 2)$	$OSp(8^* 4)$
$\ell = 0$	$B[0]_{p/2}^{[0,0,p,0]}$	$B\bar{B}[0, 0]_p^{[0,p,0]}$	$D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 2$	$Q^4 B[0]_{p/2}^{[0,0,p,0]}$	$Q^2 \bar{Q}^2 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^4 D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 0$	$Q^8 B[0]_{p/2}^{[0,0,p,0]}$	$Q^4 \bar{Q}^4 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^8 D[0, 0, 0]_{2p}^{[p,0]}$

With 16 Qs, selection rule is

$$\mathcal{O}_{k_1} \times \mathcal{O}_{k_2} \supset \mathcal{O}_p, \quad p \in \{|k_{12}| + 2, |k_{12}| + 4, \dots, k_1 + k_2 - 2\}.$$

# Operator content

Instead of  $\mathcal{M}_{\mathcal{O}}(s, t)$ , use  $\mathcal{S}_{\mathcal{O}}(s, t; \alpha)$  which is a linear combination of  $\mathcal{Y}_j(\alpha)\mathcal{M}_{\tau, \ell}(s, t)$  for all components of the superconformal block.

	$OSp(8 4)$	$PSU(2, 2 2)$	$OSp(8^* 4)$
$\ell = 0$	$B[0]_{p/2}^{[0,0,p,0]}$	$B\bar{B}[0, 0]_p^{[0,p,0]}$	$D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 2$	$Q^4 B[0]_{p/2}^{[0,0,p,0]}$	$Q^2 \bar{Q}^2 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^4 D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 0$	$Q^8 B[0]_{p/2}^{[0,0,p,0]}$	$Q^4 \bar{Q}^4 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^8 D[0, 0, 0]_{2p}^{[p,0]}$

With 16 Qs, selection rule is

$$\mathcal{O}_{k_1} \times \mathcal{O}_{k_2} \supset \mathcal{O}_p, \quad p \in \{|k_{12}| + 2, |k_{12}| + 4, \dots, k_1 + k_2 - 2\}.$$

Generic non-maximal theories can be messier [\[Dolan, Osborn; heo-th/0209056\]](#).

$$B\bar{B}[0, 0]_4^{[0,4,0]} = B\bar{B}[0, 0]_4^{2,2} \oplus A_2 \bar{A}_2[0, 0]_4^{1,1} \oplus L\bar{L}[0, 0]_4^{0,0} \oplus \dots$$

# Operator content

Instead of  $\mathcal{M}_{\mathcal{O}}(s, t)$ , use  $\mathcal{S}_{\mathcal{O}}(s, t; \alpha)$  which is a linear combination of  $\mathcal{Y}_j(\alpha)\mathcal{M}_{\tau, \ell}(s, t)$  for all components of the superconformal block.

	$OSp(8 4)$	$PSU(2, 2 2)$	$OSp(8^* 4)$
$\ell = 0$	$B[0]_{p/2}^{[0,0,p,0]}$	$B\bar{B}[0, 0]_p^{[0,p,0]}$	$D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 2$	$Q^4 B[0]_{p/2}^{[0,0,p,0]}$	$Q^2 \bar{Q}^2 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^4 D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 0$	$Q^8 B[0]_{p/2}^{[0,0,p,0]}$	$Q^4 \bar{Q}^4 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^8 D[0, 0, 0]_{2p}^{[p,0]}$

With 16 Qs, selection rule is

$$\mathcal{O}_{k_1} \times \mathcal{O}_{k_2} \supset \mathcal{O}_p, \quad p \in \{|k_{12}| + 2, |k_{12}| + 4, \dots, k_1 + k_2 - 2\}.$$

Generic non-maximal theories can be messier [\[Dolan, Osborn; heo-th/0209056\]](#).

$$B\bar{B}[0, 0]_4^{[0,4,0]} = B\bar{B}[0, 0]_4^{2,2} \oplus A_2 \bar{A}_2[0, 0]_4^{1,1} \oplus L\bar{L}[0, 0]_4^{0,0} \oplus \dots$$

If SUSY is broken with a brane, look for  $\ell = 1 \Rightarrow$  half-BPS again!



# Operator content

Instead of  $\mathcal{M}_{\mathcal{O}}(s, t)$ , use  $\mathcal{S}_{\mathcal{O}}(s, t; \alpha)$  which is a linear combination of  $\mathcal{Y}_j(\alpha)\mathcal{M}_{\tau, \ell}(s, t)$  for all components of the superconformal block.

	$OSp(8 4)$	$PSU(2, 2 2)$	$OSp(8^* 4)$
$\ell = 0$	$B[0]_{p/2}^{[0,0,p,0]}$	$B\bar{B}[0, 0]_p^{[0,p,0]}$	$D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 2$	$Q^4 B[0]_{p/2}^{[0,0,p,0]}$	$Q^2 \bar{Q}^2 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^4 D[0, 0, 0]_{2p}^{[p,0]}$
$\ell = 0$	$Q^8 B[0]_{p/2}^{[0,0,p,0]}$	$Q^4 \bar{Q}^4 B\bar{B}[0, 0]_p^{[0,p,0]}$	$Q^8 D[0, 0, 0]_{2p}^{[p,0]}$

With 16 Qs and 8 Qs, selection rule is

$$\mathcal{O}_{k_1} \times \mathcal{O}_{k_2} \supset \mathcal{O}_p, \quad p \in \{|k_{12}| + 2, |k_{12}| + 4, \dots, k_1 + k_2 - 2\}.$$

Generic non-maximal theories can be messier [\[Dolan, Osborn; heo-th/0209056\]](#).

$$B\bar{B}[0, 0]_4^{[0,4,0]} = B\bar{B}[0, 0]_4^{2,2} \oplus A_2 \bar{A}_2[0, 0]_4^{1,1} \oplus L\bar{L}[0, 0]_4^{0,0} \oplus \dots$$


If SUSY is broken with a brane, look for  $\ell = 1 \Rightarrow$  half-BPS again!

# Super-gravitons and super-gluons


Probe branes wrap  $AdS_{d+1} \times S^3$  to produce  $SU(2)_L \times SU(2)_R$ .

# Super-gravitons and super-gluons

Probe branes wrap  $AdS_{d+1} \times S^3$  to produce  $SU(2)_L \times SU(2)_R$ .




D3-D3 open strings



Closed strings


# Super-gravitons and super-gluons

Probe branes wrap  $AdS_{d+1} \times S^3$  to produce  $SU(2)_L \times SU(2)_R$ .



D3-D3 open strings

D3-D7 / D7-D3 open strings




Closed strings

D7-D7 open strings


# Super-gravitons and super-gluons

Probe branes wrap  $AdS_{d+1} \times S^3$  to produce  $SU(2)_L \times SU(2)_R$ .



D3-D3 open strings

D3-D7 / D7-D3 open strings



Closed strings

D7-D7 open strings

$$\mathcal{M}_{16Qs}(s, t; \alpha, \alpha') = \sum_{p=|k_{12}|+2}^{k_1+k_2-2} C_{12p} C_{34p} \mathcal{S}_p(s, t; \alpha, \alpha') + \text{crossed} + \text{contact}$$

$$\mathcal{M}_{8Qs}^{l_1 l_2 l_3 l_4}(s, t; \alpha, \beta) = f^{l_1 l_2 J} f^{J l_3 l_4} \sum_{p=|k_{12}|+2}^{k_1+k_2-2} C_{12p} C_{34p} \mathcal{S}_p(s, t; \alpha) \mathcal{Y}_{p-2}(\beta) + \text{same}$$

Scatter  $\mathcal{O}_k(x, t)$  at  $O(1/c_T)$  and  $\mathcal{O}'_k(x, v, \bar{v})$  at  $O(1/c_J)$ .

$$C_{k_1, k_2, k_3} = ?$$

Determined in terms of  $\Xi = \frac{1}{2} \sum_i k_i$  and  $\alpha_i = \Xi - k_i$ .

$$C_{k_1, k_2, k_3} = ?$$

Determined in terms of  $\Xi = \frac{1}{2} \sum_i k_i$  and  $\alpha_i = \Xi - k_i$ .

$$C_{k_1, k_2, k_3}^{16Qs} = \begin{cases} \sqrt{\frac{70}{\pi^3 c_T}} 2^{2\Xi-1} \Gamma[\Xi] \prod_i \frac{\Gamma[\alpha_i + \frac{1}{2}]}{\sqrt{\Gamma[2k_i - 1]}} & AdS_7 \\ \sqrt{\frac{30k_1 k_2 k_3}{c_T}} & AdS_5 \\ \sqrt{\frac{\pi}{3c_T}} \frac{2^{3-\Xi}}{\Gamma[\frac{\Xi+2}{2}]} \prod_i \frac{\sqrt{\Gamma[k_i+2]}}{\Gamma[\frac{\alpha_i+1}{2}]} & AdS_4 \end{cases}$$

from [Lee, Minwalla, Rangamani, Seiberg; hep-th/9806074] [Bastianelli, Zucchini; hep-th/9907047] .

$$C_{k_1, k_2, k_3} = ?$$

Determined in terms of  $\Xi = \frac{1}{2} \sum_i k_i$  and  $\alpha_i = \Xi - k_i$ .

$$C_{k_1, k_2, k_3}^{16Qs} = \begin{cases} \sqrt{\frac{70}{\pi^3 c_T}} 2^{2\Xi-1} \Gamma[\Xi] \prod_i \frac{\Gamma[\alpha_i + \frac{1}{2}]}{\sqrt{\Gamma[2k_i - 1]}} & AdS_7 \\ \sqrt{\frac{30 k_1 k_2 k_3}{c_T}} & AdS_5 \\ \sqrt{\frac{\pi}{3 c_T}} \frac{2^{3-\Xi}}{\Gamma[\frac{\Xi+2}{2}]} \prod_i \frac{\sqrt{\Gamma[k_i+2]}}{\Gamma[\frac{\alpha_i+1}{2}]} & AdS_4 \end{cases}$$

from [Lee, Minwalla, Rangamani, Seiberg; hep-th/9806074] [Bastianelli, Zucchini; hep-th/9907047] .

$$C_{k_1, k_2, k_3}^{8Qs} = \begin{cases} \sqrt{\frac{30}{\pi c_J}} \Gamma[\Xi - \frac{1}{2}] \prod_i \frac{\Gamma[\alpha_i + \frac{1}{2}]}{\sqrt{\Gamma[k_i \pm \frac{1}{2}]}} & AdS_7 \\ \frac{2^{2\Xi+23} 7^{-\frac{6\Xi}{4}} \Gamma[\frac{\Xi \pm 1}{2}]}{\sqrt{3\pi c_J} \Gamma[\frac{\Xi}{2}]} \prod_i \sqrt{\frac{\Gamma[k_i]}{\Gamma[k_i \pm \frac{1}{3}]}} \frac{\Gamma[\frac{3}{2}\alpha_i]}{\Gamma[\alpha_i]} & AdS_6 \\ \sqrt{\frac{6}{c_J}} & AdS_5 \\ \sqrt{\frac{2\pi}{c_J}} \frac{2^{3-\Xi}}{\Gamma[\frac{\Xi}{2}]} \prod_i \frac{\sqrt{\Gamma[k_i]}}{\Gamma[\frac{\alpha_i+1}{2}]} & AdS_4 \text{ (really 6Qs)} \end{cases}$$

from [Alday, CB, Ferrero, Zhou; 2103.15830] .



# 1. Superconformal Ward identity

Apply  $(\chi\partial_\chi - \epsilon\alpha\partial_\alpha)\mathcal{G}|_{\alpha=\chi^{-1}}=0$  in Mellin space [\[Zhou; 1712.02800\]](#).

$$U\partial_U \rightarrow \left[ \frac{s}{2} - \epsilon \frac{k_1 + k_2 - 2\mathcal{E}}{2} \right] \times, \quad V\partial_V \rightarrow \left[ \frac{t}{2} + \epsilon \frac{k_1 - k_4 - 2\mathcal{E}}{2} \right] \times$$
$$U^m V^n \circ \mathcal{M}(s, t) = \left( \frac{\Delta_1 + \Delta_2 - s}{2} \right)_m \left( \frac{\Delta_3 + \Delta_4 - s}{2} \right)_m \left( \frac{\Delta_1 + \Delta_4 - t}{2} \right)_n$$
$$\left( \frac{\Delta_2 + \Delta_3 - t}{2} \right)_n \left( \frac{\Delta_1 + \Delta_3 - u}{2} \right)_{-m-n} \left( \frac{\Delta_2 + \Delta_4 - u}{2} \right)_{-m-n} \mathcal{M}(s - 2m, t - 2n)$$

# 1. Superconformal Ward identity

Apply  $(\chi\partial_\chi - \epsilon\alpha\partial_\alpha)\mathcal{G}|_{\alpha=\chi^{-1}}=0$  in Mellin space [Zhou; 1712.02800].

$$U\partial_U \rightarrow \left[ \frac{s}{2} - \epsilon \frac{k_1 + k_2 - 2\mathcal{E}}{2} \right] \times, \quad V\partial_V \rightarrow \left[ \frac{t}{2} + \epsilon \frac{k_1 - k_4 - 2\mathcal{E}}{2} \right] \times$$
$$U^m V^n \circ \mathcal{M}(s, t) = \left( \frac{\Delta_1 + \Delta_2 - s}{2} \right)_m \left( \frac{\Delta_3 + \Delta_4 - s}{2} \right)_m \left( \frac{\Delta_1 + \Delta_4 - t}{2} \right)_n$$
$$\left( \frac{\Delta_2 + \Delta_3 - t}{2} \right)_n \left( \frac{\Delta_1 + \Delta_3 - u}{2} \right)_{-m-n} \left( \frac{\Delta_2 + \Delta_4 - u}{2} \right)_{-m-n} \mathcal{M}(s - 2m, t - 2n)$$

Residues of  $\mathcal{S}_p$  can have  $m$  dependence simplified.

$$\frac{\sigma^i \tau^j H_{p,m}^{i,j}}{s - \epsilon p - 2m} [t^2 + q_1^p(m)t + q_2^p(m)] \quad \text{or} \quad \frac{(1 - \alpha)^i H_{p,m}^i}{s - \epsilon p - 2m} [t + q_1^p(m)]$$
$$s + t + u = \epsilon \sum_i k_i, \quad s = \tau + 2m$$

# 1. Superconformal Ward identity

Apply  $(\chi\partial_\chi - \epsilon\alpha\partial_\alpha)\mathcal{G}|_{\alpha=\chi^{-1}}=0$  in Mellin space [Zhou; 1712.02800].

$$U\partial_U \rightarrow \left[ \frac{s}{2} - \epsilon \frac{k_1 + k_2 - 2\mathcal{E}}{2} \right] \times, \quad V\partial_V \rightarrow \left[ \frac{t}{2} + \epsilon \frac{k_1 - k_4 - 2\mathcal{E}}{2} \right] \times$$
$$U^m V^n \circ \mathcal{M}(s, t) = \left( \frac{\Delta_1 + \Delta_2 - s}{2} \right)_m \left( \frac{\Delta_3 + \Delta_4 - s}{2} \right)_m \left( \frac{\Delta_1 + \Delta_4 - t}{2} \right)_n$$
$$\left( \frac{\Delta_2 + \Delta_3 - t}{2} \right)_n \left( \frac{\Delta_1 + \Delta_3 - u}{2} \right)_{-m-n} \left( \frac{\Delta_2 + \Delta_4 - u}{2} \right)_{-m-n} \mathcal{M}(s - 2m, t - 2n)$$

Residues of  $\mathcal{S}_p$  can have  $m$  dependence simplified.

$$\frac{\sigma^i \tau^j H_{p,m}^{i,j}}{s - \epsilon p - 2m} [t^2 + q_1^p(m)t + q_2^p(m)] \quad \text{or} \quad \frac{(1 - \alpha)^i H_{p,m}^i}{s - \epsilon p - 2m} [t + q_1^p(m)]$$
$$s + t + u = \epsilon \sum_i k_i, \quad s = \tau + 2m$$

Leads to amplitudes with **no** additional contact terms [Alday, Zhou; 2006.06653].

## 2. Flat space limit

Amplitudes take a universal form for  $s, t \rightarrow \infty$  with  $s + t + u = 0$ .

$$\frac{\mathcal{M}_{16Q_5}(s, t, \alpha, \alpha')}{P_{\mathcal{E}-2}(\sigma, \tau)} = \frac{(s + t - \alpha s)^2 (s + t - \alpha' s)^2}{stu}$$

## 2. Flat space limit

Amplitudes take a universal form for  $s, t \rightarrow \infty$  with  $s + t + u = 0$ .

$$\frac{\mathcal{M}_{16Q_5}(s, t, \alpha, \alpha')}{P_{\mathcal{E}-2}(\sigma, \tau)} = \frac{(s + t - \alpha s)^2 (s + t - \alpha' s)^2}{stu}$$

With color and kinematic factors

$$c_s = f^{h_1 l_2 J} f^{J l_3 l_4}, \quad c_t = f^{h_1 l_4 J} f^{J l_2 l_3}, \quad c_u = f^{h_1 l_3 J} f^{J l_4 l_2}$$

$$N_s = u(1 - \alpha) - t\alpha, \quad N_t = (\alpha - 1)(u + s\alpha), \quad N_u = \alpha(t + s(1 - \alpha))$$

gluon analogue agrees with [\[Adamo, Casali, Mason, Nekovar; 1810.05115\]](#) :

$$\frac{\mathcal{M}_{8Q_5}(s, t, \alpha, \beta)}{P_{\mathcal{E}-2}(\sigma, \tau)} = \frac{(tc_s - sc_t)(s + t - \alpha s)^2}{stu} = \left[ \frac{c_s N_s}{s} + \frac{c_t N_t}{t} + \frac{c_u N_u}{u} \right].$$

## 2. Flat space limit

Amplitudes take a universal form for  $s, t \rightarrow \infty$  with  $s + t + u = 0$ .

$$\frac{\mathcal{M}_{16Qs}(s, t, \alpha, \alpha')}{P_{\mathcal{E}-2}(\sigma, \tau)} = \frac{(s + t - \alpha s)^2 (s + t - \alpha' s)^2}{stu}$$

With color and kinematic factors

$$c_s = f^{h_1 l_2 J} f^{J l_3 l_4}, \quad c_t = f^{h_1 l_4 J} f^{J l_2 l_3}, \quad c_u = f^{h_1 l_3 J} f^{J l_4 l_2}$$

$$N_s = u(1 - \alpha) - t\alpha, \quad N_t = (\alpha - 1)(u + s\alpha), \quad N_u = \alpha(t + s(1 - \alpha))$$

gluon analogue agrees with [\[Adamo, Casali, Mason, Nekovar; 1810.05115\]](#) :

$$\frac{\mathcal{M}_{8Qs}(s, t, \alpha, \beta)}{P_{\mathcal{E}-2}(\sigma, \tau)} = \frac{(tc_s - sc_t)(s + t - \alpha s)^2}{stu} = \left[ \frac{c_s N_s}{s} + \frac{c_t N_t}{t} + \frac{c_u N_u}{u} \right].$$

Polynomial given by the overlap of wavefunctions on a transverse  $S^d$ .

$$P_{\mathcal{E}-2}(\sigma, \tau) \propto \int_{S^d} dT (t_1 \cdot T)^{k_1-2} (t_2 \cdot T)^{k_2-2} (t_3 \cdot T)^{k_3-2} (t_4 \cdot T)^{k_4-2}$$

### 3. Chiral algebra

Superconformal Ward identities in four dimensions are

$$\partial_{\chi'} \mathcal{G}^{\mathcal{N}=2}(\chi, \chi'; \chi'^{-1}) = 0, \quad \partial_{\chi'} \mathcal{G}^{\mathcal{N}=4}(\chi, \chi'; \alpha, \chi'^{-1}) = 0.$$

Solutions have the form  $\mathcal{G} = \mathcal{K} + R\mathcal{H}$  with

$$R^{\mathcal{N}=2} = (1 - \alpha\chi)(1 - \alpha\chi'), \quad R^{\mathcal{N}=4} = (1 - \alpha\chi)(1 - \alpha\chi')(1 - \alpha'\chi)(1 - \alpha'\chi').$$

Cohomological explanation: [\[Beem, Lemos, Liendo, Peelaers, Rastelli, van Rees; 1312.5344\]](#) .

### 3. Chiral algebra


Superconformal Ward identities in four dimensions are

$$\partial_{\chi'} \mathcal{G}^{\mathcal{N}=2}(\chi, \chi'; \chi'^{-1}) = 0, \quad \partial_{\chi'} \mathcal{G}^{\mathcal{N}=4}(\chi, \chi'; \alpha, \chi'^{-1}) = 0.$$

Solutions have the form  $\mathcal{G} = \mathcal{K} + R\mathcal{H}$  with

$$R^{\mathcal{N}=2} = (1 - \alpha\chi)(1 - \alpha\chi'), \quad R^{\mathcal{N}=4} = (1 - \alpha\chi)(1 - \alpha\chi')(1 - \alpha'\chi)(1 - \alpha'\chi').$$

Cohomological explanation: [\[Beem, Lemos, Liendo, Peelaers, Rastelli, van Rees; 1312.5344\]](#) .



Plane preserved by  $\mathfrak{sl}(2) \times \mathfrak{sl}(2|2)$  admits nilpotent  $\mathbb{Q}$ .

R symmetry gives  $\bar{L}_{0,\pm 1} - R_{0,\pm 1} = \{\mathbb{Q}, \cdot\}$  commuting with  $L_{0,\pm 1}$ .

2d and 4d central charges related by a negative factor.



### 3. Chiral algebra

Bootstrapping a W-algebra means solving for  $\lambda_{12\mathcal{O}}$ .

$$\mathcal{O}_1(z)\mathcal{O}_2(0) = \sum_{\mathcal{O}} \lambda_{12\mathcal{O}} \sum_{m=0}^{\infty} \frac{(h+h_{12})_m}{m!(2h)_m} \frac{\partial^m \mathcal{O}(0)}{z^{h_1+h_2-h-m}}$$

### 3. Chiral algebra

Bootstrapping a W-algebra means solving for  $\lambda_{12\mathcal{O}}$ .

$$\mathcal{O}_1(z)\mathcal{O}_2(0) = \sum_{\mathcal{O}} \lambda_{12\mathcal{O}} \sum_{m=0}^{\infty} \frac{(h+h_{12})_m}{m!(2h)_m} \frac{\partial^m \mathcal{O}(0)}{z^{h_1+h_2-h-m}}$$

Crossing under  $1 \leftrightarrow 3$  manifest but  $1 \leftrightarrow 4$  must **still be imposed**.

$$F_{1234}(\chi) + \frac{(-1)^{k_1+k_4} \chi^{k_1+k_2}}{(\chi-1)^{k_2+k_3}} F_{3214}(1-\chi)$$

$$F_{1234}(\chi) = \sum_p C_{12p} C_{34p} \sum_{m=0}^{k_1+k_2-p-1} \frac{(p-k_{12})_m (p+k_{34})_m}{m!(2p)_m \chi^{k_1+k_2-p-m}}$$

### 3. Chiral algebra

Bootstrapping a W-algebra means solving for  $\lambda_{12\mathcal{O}}$ .

$$\mathcal{O}_1(z)\mathcal{O}_2(0) = \sum_{\mathcal{O}} \lambda_{12\mathcal{O}} \sum_{m=0}^{\infty} \frac{(h+h_{12})_m}{m!(2h)_m} \frac{\partial^m \mathcal{O}(0)}{z^{h_1+h_2-h-m}}$$

Crossing under  $1 \leftrightarrow 3$  manifest but  $1 \leftrightarrow 4$  must **still be imposed**.

$$F_{1234}(\chi) + \frac{(-1)^{k_1+k_4} \chi^{k_1+k_2}}{(\chi-1)^{k_2+k_3}} F_{3214}(1-\chi)$$

$$F_{1234}(\chi) = \sum_p C_{12p} C_{34p} \sum_{m=0}^{k_1+k_2-p-1} \frac{(p-k_{12})_m (p+k_{34})_m}{m!(2p)_m \chi^{k_1+k_2-p-m}}$$

All OPE coefficients fixed in Argyres-Douglas case.

$$F_{1234}^{l_1 l_2 l_3 l_4}(\chi; \beta) = \sqrt{\frac{6}{c_J}} f^{l_1 l_2 J} f^{J l_3 l_4} \sum_p g_{1-\frac{p}{2}}^{\frac{k_{21}}{2}, \frac{k_{43}}{2}} (\beta^{-1})^{\frac{k_1+k_2-p-2}{2}} \sum_{m=0}^{\frac{(p-k_{12})_m (p+k_{34})_m}{m!(p)_m \chi^{\frac{k_1+k_2-p}{2}-m}}$$

### 3. Chiral algebra and loops

Simplest chiral algebra correlator in 4d  $\mathcal{N} = 4$  SYM.

$$F_{2222}(\chi; \alpha) = 1 + (\chi\alpha)^2 + \chi^2 \left( \frac{\alpha - 1}{1 - \chi} \right)^2 + \frac{12}{c} \left[ \frac{\chi}{1 - \chi} - 2\chi\alpha + \frac{(\chi\alpha)^2}{1 - \chi} \right]$$

Why do we have  $F_{2222}(\chi; \alpha)|_{1/c^2} = 0$  but  $F_{3333}(\chi; \alpha)|_{1/c^2} \neq 0$ ?

### 3. Chiral algebra and loops

Simplest chiral algebra correlator in 4d  $\mathcal{N} = 4$  SYM.

$$F_{2222}(\chi; \alpha) = 1 + (\chi\alpha)^2 + \chi^2 \left( \frac{\alpha - 1}{1 - \chi} \right)^2 + \frac{12}{c} \left[ \frac{\chi}{1 - \chi} - 2\chi\alpha + \frac{(\chi\alpha)^2}{1 - \chi} \right]$$

Why do we have  $F_{2222}(\chi; \alpha)|_{1/c^2} = 0$  but  $F_{3333}(\chi; \alpha)|_{1/c^2} \neq 0$ ?

$$J^{(2)}(z)J^{(2)}(0) \sim \frac{1}{z^2} + \frac{J^{(2)}(0)}{\sqrt{c}z} + \text{regular}$$

$$J^{(3)}(z)J^{(3)}(0) \sim \frac{1}{z^3} + \frac{J^{(2)}(0)}{\sqrt{c}z^2} + \frac{J^{(4)}(0)}{\sqrt{c}z} + \text{regular}$$

First OPE **cannot** pick up any  $\frac{1}{c}J^{(a)}J^{(b)}$  but second OPE **can** pick up  $\frac{1}{c}J^{(2)}J^{(2)}$  at order  $\frac{1}{z}$ .

### 3. Chiral algebra and loops

Simplest chiral algebra correlator in 4d  $\mathcal{N} = 4$  SYM.

$$F_{2222}(\chi; \alpha) = 1 + (\chi\alpha)^2 + \chi^2 \left( \frac{\alpha - 1}{1 - \chi} \right)^2 + \frac{12}{c} \left[ \frac{\chi}{1 - \chi} - 2\chi\alpha + \frac{(\chi\alpha)^2}{1 - \chi} \right]$$

Why do we have  $F_{2222}(\chi; \alpha)|_{1/c^2} = 0$  but  $F_{3333}(\chi; \alpha)|_{1/c^2} \neq 0$ ?

$$J^{(2)}(z)J^{(2)}(0) \sim \frac{1}{z^2} + \frac{J^{(2)}(0)}{\sqrt{cz}} + \text{regular}$$

$$J^{(3)}(z)J^{(3)}(0) \sim \frac{1}{z^3} + \frac{J^{(2)}(0)}{\sqrt{cz^2}} + \frac{J^{(4)}(0)}{\sqrt{cz}} + \text{regular}$$

First OPE **cannot** pick up any  $\frac{1}{c}J^{(a)}J^{(b)}$  but second OPE **can** pick up  $\frac{1}{c}J^{(2)}J^{(2)}$  at order  $\frac{1}{z}$ .

$$F_{3333}(\chi; \alpha)|_{1/c^2} = \frac{2025\chi^2\alpha(1 - \alpha)}{c^2(1 - \chi)}$$

Crossing continues to fix loops [\[CB, Ferrero, Zhou; 2101.04114\]](#).

# Unprotected data at one loop

For  $[\mathcal{O}_2\mathcal{O}_2]_{n,\ell} \subset \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2 \rangle$ , it is standard to use  $\mathcal{H}(\chi, \chi')$ .

$$\mathcal{H}(\chi, \chi') = \sum_{n,\ell} a_{n,\ell}^{(0)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi')$$

# Unprotected data at one loop

For  $[\mathcal{O}_2\mathcal{O}_2]_{n,\ell} \subset \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2 \rangle$ , it is standard to use  $\mathcal{H}(\chi, \chi')$ .

$$\mathcal{H}(\chi, \chi') = \sum_{n,\ell} \left( a_{n,\ell}^{(0)} + a_{n,\ell}^{(1)} + \dots \right) G_{\Delta_{n,\ell} + \gamma_{n,\ell}^{(1)} + \dots, \ell}(\chi, \chi')$$



# Unprotected data at one loop

For  $[\mathcal{O}_2\mathcal{O}_2]_{n,\ell} \subset \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2 \rangle$ , it is standard to use  $\mathcal{H}(\chi, \chi')$ .

$$\begin{aligned}\mathcal{H}(\chi, \chi') &= \sum_{n,\ell} a_{n,\ell}^{(0)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \sum_{n,\ell} \left[ a_{n,\ell}^{(1)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \gamma_{n,\ell}^{(1)} a_{n,\ell}^{(0)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right] \\ &+ \sum_{n,\ell} \left[ a_{n,\ell}^{(2)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + 2\gamma_{n,\ell}^{(1)} a_{n,\ell}^{(1)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \gamma_{n,\ell}^{(2)} a_{n,\ell}^{(0)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right. \\ &\left. + \gamma_{n,\ell}^{(1)2} a_{n,\ell}^{(0)} G''_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right] + O(1/c_T^3)\end{aligned}$$

# Unprotected data at one loop

For  $[\mathcal{O}_2\mathcal{O}_2]_{n,\ell} \subset \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2 \rangle$ , it is standard to use  $\mathcal{H}(\chi, \chi')$ .

$$\begin{aligned}\mathcal{H}(\chi, \chi') &= \sum_{n,\ell} a_{n,\ell}^{(0)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \sum_{n,\ell} \left[ a_{n,\ell}^{(1)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \gamma_{n,\ell}^{(1)} a_{n,\ell}^{(0)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right] \\ &+ \sum_{n,\ell} \left[ a_{n,\ell}^{(2)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + 2\gamma_{n,\ell}^{(1)} a_{n,\ell}^{(1)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \gamma_{n,\ell}^{(2)} a_{n,\ell}^{(0)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right. \\ &\left. + \gamma_{n,\ell}^{(1)2} a_{n,\ell}^{(0)} G''_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right] + O(1/c_T^3)\end{aligned}$$

Double-log is also a **double-discontinuity** defined by

$$dDisc[f(\chi, \chi')] = f(\chi, \chi') - \frac{1}{2} [f(\chi, e^{2\pi i} \chi') + f(\chi, e^{-2\pi i} \chi')].$$

# Unprotected data at one loop

For  $[\mathcal{O}_2\mathcal{O}_2]_{n,\ell} \subset \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2 \rangle$ , it is standard to use  $\mathcal{H}(\chi, \chi')$ .

$$\begin{aligned}\mathcal{H}(\chi, \chi') &= \sum_{n,\ell} a_{n,\ell}^{(0)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \sum_{n,\ell} \left[ a_{n,\ell}^{(1)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \gamma_{n,\ell}^{(1)} a_{n,\ell}^{(0)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right] \\ &+ \sum_{n,\ell} \left[ a_{n,\ell}^{(2)} G_{\Delta_{n,\ell,\ell}}(\chi, \chi') + 2\gamma_{n,\ell}^{(1)} a_{n,\ell}^{(1)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') + \gamma_{n,\ell}^{(2)} a_{n,\ell}^{(0)} G'_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right. \\ &\left. + \gamma_{n,\ell}^{(1)2} a_{n,\ell}^{(0)} G''_{\Delta_{n,\ell,\ell}}(\chi, \chi') \right] + O(1/c_T^3)\end{aligned}$$

Double-log is also a **double-discontinuity** defined by


$$dDisc[f(\chi, \chi')] = f(\chi, \chi') - \frac{1}{2} [f(\chi, e^{2\pi i} \chi') + f(\chi, e^{-2\pi i} \chi')].$$

Full spectral density encoded in this [\[Caron-Huot; 1703.00278\]](#) !

$$c(\Delta, \ell) = \kappa_{\Delta,\ell} \int_0^1 \int_0^1 \left| \frac{\chi - \chi'}{\chi\chi'} \right|^{d-2} G_{\ell+d-1, \Delta+1-d}(\chi, \chi') dDisc[\mathcal{H}(\chi, \chi')] \frac{d\chi}{\chi^2} \frac{d\chi'}{\chi'^2}$$


# Unprotected data at one loop

Use  $\langle \mathcal{O}_2 \mathcal{O}_2 \mathcal{O}_p \mathcal{O}_p \rangle$  to get around  $\langle \gamma^{(1)2} a^{(0)} \rangle_{n,\ell} \neq \frac{\langle \gamma^{(1)} a^{(0)} \rangle_{n,\ell}^2}{\langle a^{(0)} \rangle_{n,\ell}}$ .



# Unprotected data at one loop

Use  $\langle \mathcal{O}_2 \mathcal{O}_2 \mathcal{O}_p \mathcal{O}_p \rangle$  to get around  $\langle \gamma^{(1)2} a^{(0)} \rangle_{n,\ell} \neq \frac{\langle \gamma^{(1)} a^{(0)} \rangle_{n,\ell}^2}{\langle a^{(0)} \rangle_{n,\ell}}$ .




$$\begin{pmatrix} [\mathcal{O}_2 \mathcal{O}_2]_1 \\ [\mathcal{O}_3 \mathcal{O}_3]_0 \end{pmatrix} = \begin{pmatrix} \frac{\lambda_{22A}}{\sqrt{\lambda_{22A}^2 + \lambda_{22B}^2}} & \frac{\lambda_{22B}}{\sqrt{\lambda_{22A}^2 + \lambda_{22B}^2}} \\ \frac{\lambda_{33A}}{\sqrt{\lambda_{33A}^2 + \lambda_{33B}^2}} & \frac{\lambda_{33B}}{\sqrt{\lambda_{33A}^2 + \lambda_{33B}^2}} \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix}$$

$$M = Q \begin{pmatrix} \gamma_A^{(1)} & 0 \\ 0 & \gamma_B^{(1)} \end{pmatrix} Q^T, \quad \langle \gamma^{(1)2} a^{(0)} \rangle_1 \in M^2$$

# Unprotected data at one loop

Use  $\langle \mathcal{O}_2 \mathcal{O}_2 \mathcal{O}_p \mathcal{O}_p \rangle$  to get around  $\langle \gamma^{(1)2} a^{(0)} \rangle_{n,\ell} \neq \frac{\langle \gamma^{(1)} a^{(0)} \rangle_{n,\ell}^2}{\langle a^{(0)} \rangle_{n,\ell}}$ .



$$\begin{pmatrix} [\mathcal{O}_2 \mathcal{O}_2]_1 \\ [\mathcal{O}_3 \mathcal{O}_3]_0 \end{pmatrix} = \begin{pmatrix} \frac{\lambda_{22A}}{\sqrt{\lambda_{22A}^2 + \lambda_{22B}^2}} & \frac{\lambda_{22B}}{\sqrt{\lambda_{22A}^2 + \lambda_{22B}^2}} \\ \frac{\lambda_{33A}}{\sqrt{\lambda_{33A}^2 + \lambda_{33B}^2}} & \frac{\lambda_{33B}}{\sqrt{\lambda_{33A}^2 + \lambda_{33B}^2}} \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix}$$

$$M = Q \begin{pmatrix} \gamma_A^{(1)} & 0 \\ 0 & \gamma_B^{(1)} \end{pmatrix} Q^T, \quad \langle \gamma^{(1)2} a^{(0)} \rangle_1 \in M^2$$

Eigenvalues never involve higher roots [\[Aprile, Drummond, Heslop, Paul; 1706.02822\]](#) !

$$\gamma_{n,\ell,i}^{(1)} = -\frac{2(n+1)_4(n+\ell+2)_4}{(\ell+2i+1)_6}, \quad i = 0, \dots, n$$

# Hidden conformal symmetry

Higher dimensional blocks diagonalize this [\[Caron-Huot, Trinh; 1809.09173\]](#) .

# Hidden conformal symmetry

Higher dimensional blocks diagonalize this [\[Caron-Huot, Trinh; 1809.09173\]](#).

$$\begin{aligned}\widetilde{\mathcal{M}}_{2222}(s, t) &\rightarrow \mathcal{A}_{grav}(s, t) \propto \frac{1}{stu} \\ \widetilde{\mathcal{M}}_{2222}^{h_1 l_2 l_3 l_4}(s, t) &\rightarrow \mathcal{A}_{glu}^{h_1 l_2 l_3 l_4}(s, t) \propto \frac{f^{h_1 l_2 J} f^{J l_3 l_4}}{su} - \frac{f^{h_1 l_4 J} f^{J l_2 l_3}}{tu}\end{aligned}$$

Graviton (gluon) amplitude conformal for  $d = 10$  ( $d = 8$ )!

$$K_\mu = \sum_{i=1}^4 \left[ \frac{p_{i\mu}}{2} \frac{\partial}{\partial p_i} \cdot \frac{\partial}{\partial p_i} - p_i^\nu \frac{\partial}{\partial p_i^\nu} \frac{\partial}{\partial p_i^\mu} - \frac{d-2}{2} \frac{\partial}{\partial p_i^\mu} \right]$$



# Hidden conformal symmetry

Higher dimensional blocks diagonalize this [\[Caron-Huot, Trinh; 1809.09173\]](#).

$$\begin{aligned}\widetilde{\mathcal{M}}_{2222}(s, t) &\rightarrow \mathcal{A}_{grav}(s, t) \propto \frac{1}{stu} \\ \widetilde{\mathcal{M}}_{2222}^{h_1 l_2 l_3 l_4}(s, t) &\rightarrow \mathcal{A}_{glu}^{h_1 l_2 l_3 l_4}(s, t) \propto \frac{f^{h_1 l_2 J} f^{J l_3 l_4}}{su} - \frac{f^{h_1 l_4 J} f^{J l_2 l_3}}{tu}\end{aligned}$$

Graviton (gluon) amplitude conformal for  $d = 10$  ( $d = 8$ )!

$$K_\mu = \sum_{i=1}^4 \left[ \frac{p_{i\mu}}{2} \frac{\partial}{\partial p_i} \cdot \frac{\partial}{\partial p_i} - p_i^\nu \frac{\partial}{\partial p_i^\nu} \frac{\partial}{\partial p_i^\mu} - \frac{d-2}{2} \frac{\partial}{\partial p_i^\mu} \right]$$

Turns lowest KK mode into **generating function** for all others.

$$\mathcal{H}_{k_1, k_2, k_3, k_4}(x_{ij}^2) \subset \mathcal{H}_{2222}(x_{ij}^2 + t_{ij})$$

# Hidden conformal symmetry

Higher dimensional blocks diagonalize this [\[Caron-Huot, Trinh; 1809.09173\]](#).

$$\begin{aligned}\widetilde{\mathcal{M}}_{2222}(s, t) &\rightarrow \mathcal{A}_{grav}(s, t) \propto \frac{1}{stu} \\ \widetilde{\mathcal{M}}_{2222}^{h_1 l_2 l_3 l_4}(s, t) &\rightarrow \mathcal{A}_{glu}^{h_1 l_2 l_3 l_4}(s, t) \propto \frac{f^{h_1 l_2 J} f^{J l_3 l_4}}{su} - \frac{f^{h_1 l_4 J} f^{J l_2 l_3}}{tu}\end{aligned}$$

Graviton (gluon) amplitude conformal for  $d = 10$  ( $d = 8$ )!

$$K_\mu = \sum_{i=1}^4 \left[ \frac{p_{i\mu}}{2} \frac{\partial}{\partial p_i} \cdot \frac{\partial}{\partial p_i} - p_i^\nu \frac{\partial}{\partial p_i^\nu} \frac{\partial}{\partial p_i^\mu} - \frac{d-2}{2} \frac{\partial}{\partial p_i^\mu} \right]$$

Turns lowest KK mode into **generating function** for all others.

$$\mathcal{H}_{k_1, k_2, k_3, k_4}(x_{ij}^2) \subset \mathcal{H}_{2222}(x_{ij}^2 + t_{ij})$$

Background  $AdS_5 \times S^5$  and brane locus  $AdS_5 \times S^3$  are both conformally flat. Corrections in  $\lambda$  also organize this way [\[Caron-Huot, Coronado; 2106.03892\]](#).

# An old conjecture

Actions for different  $d$  are perturbatively equivalent [\[Parisi, Sourlas; 1979\]](#) .

$$S = \int d^d x d\theta d\bar{\theta} \left[ -\frac{1}{2} \Phi \partial^2 \Phi + V(\Phi) \right] \leftrightarrow S = \int d^{d-2} x \left[ -\frac{1}{2} \phi \partial^2 \phi + V(\phi) \right]$$

# An old conjecture

Actions for different  $d$  are perturbatively equivalent [Parisi, Sourlas; 1979] .

$$S = \int d^d x d\theta d\bar{\theta} \left[ -\frac{1}{2} \Phi \partial^2 \Phi + V(\Phi) \right] \leftrightarrow S = \int d^{d-2} x \left[ -\frac{1}{2} \phi \partial^2 \phi + V(\phi) \right]$$

LHS invariant under  $\mathfrak{osp}(d+1, 1|2)$  superalgebra.

$$\begin{aligned} \delta_{\mu\nu} &\rightarrow \delta_{\mu\nu} \otimes \Omega_{+-} \\ P_\mu &\rightarrow (P_\mu, Q_+, Q_-), \quad K_\mu \rightarrow (K_\mu, S_+, S_-) \\ &\text{Similar for } M_{\mu\nu} \end{aligned}$$

# An old conjecture

Actions for different  $d$  are perturbatively equivalent [Parisi, Sourlas; 1979] .

$$S = \int d^d x d\theta d\bar{\theta} \left[ -\frac{1}{2} \Phi \partial^2 \Phi + V(\Phi) \right] \leftrightarrow S = \int d^{d-2} x \left[ -\frac{1}{2} \phi \partial^2 \phi + V(\phi) \right]$$

LHS invariant under  $\mathfrak{osp}(d+1, 1|2)$  superalgebra.

$$\begin{aligned} \delta_{\mu\nu} &\rightarrow \delta_{\mu\nu} \otimes \Omega_{+-} \\ P_\mu &\rightarrow (P_\mu, Q_+, Q_-), \quad K_\mu \rightarrow (K_\mu, S_+, S_-) \\ &\text{Similar for } M_{\mu\nu} \end{aligned}$$

Superblock can be expressed in two ways [Kaviraj, Rychkov, Trevisani; 1912.01617] .

$$G_{\Delta, \ell}^{(d-2)} = G_{\Delta, \ell}^{(d)} + c_{2,0} G_{\Delta+2, \ell}^{(d)} + c_{1,-1} G_{\Delta+1, \ell-1}^{(d)} + c_{0,-2} G_{\Delta, \ell-2}^{(d)} + c_{2,-2} G_{\Delta+2, \ell-2}^{(d)}$$

Five term relation has two terms for  $\ell = 0$ .

# Parisi-Sourlas SUSY in holography

Residues of  $\mathcal{S}_p$  involve  $K_p^{i,j}(t, u)H_{p,m}^{i,j}$  or  $K_p^i(t, u)H_{p,m}^i$ .

$K_p^{i,j}(t, u)$	$\hat{K}_p$
$t + \Delta_1 - \Delta_4 - 2\epsilon\mathcal{E}$	$2V\partial_V$
$u - \Delta_1 - \Delta_3$	$-2U\partial_U - 2V\partial_V$
$i, j$	$\sigma\partial_\sigma, \tau\partial_\tau$

# Parisi-Sourlas SUSY in holography

Residues of  $\mathcal{S}_p$  involve  $K_p^{i,j}(t, u)H_{p,m}^{i,j}$  or  $K_p^i(t, u)H_{p,m}^i$ .

$K_p^{i,j}(t, u)$	$\hat{K}_p$
$t + \Delta_1 - \Delta_4 - 2\epsilon\mathcal{E}$	$2V\partial_V$
$u - \Delta_1 - \Delta_3$	$-2U\partial_U - 2V\partial_V$
$i, j$	$\sigma\partial_\sigma, \tau\partial_\tau$

Special linear combinations are acted on by  $\hat{K}_p$  [CB, Ferrero, Zhou; 2101.04114].

$$\mathcal{M}_{\epsilon p, 0}^{(d-2)} = \mathcal{M}_{\epsilon p, 0}^{(d)} + c_{2,0}^{(d)} \mathcal{M}_{\epsilon p+2, 0}^{(d)}$$

$$\mathcal{M}_{\epsilon p, 0}^{(d-4)} = \mathcal{M}_{\epsilon p, 0}^{(d-2)} + c_{2,0}^{(d-2)} \mathcal{M}_{\epsilon p+2, 0}^{(d-2)}$$

$$= \left[ \mathcal{M}_{\epsilon p, 0}^{(d)} + c_{2,0}^{(d)} \mathcal{M}_{\epsilon p+2, 0}^{(d)} \right] + c_{2,0}^{(d-2)} \left[ \mathcal{M}_{\epsilon p+2, 0}^{(d)} + c_{2,0}^{(d)} \mathcal{M}_{\epsilon p+4, 0}^{(d)} \right]$$

# Parisi-Sourlas SUSY in holography

Residues of  $\mathcal{S}_p$  involve  $K_p^{i,j}(t, u)H_{p,m}^{i,j}$  or  $K_p^i(t, u)H_{p,m}^i$ .

$K_p^{i,j}(t, u)$	$\hat{K}_p$
$t + \Delta_1 - \Delta_4 - 2\epsilon\mathcal{E}$	$2V\partial_V$
$u - \Delta_1 - \Delta_3$	$-2U\partial_U - 2V\partial_V$
$i, j$	$\sigma\partial_\sigma, \tau\partial_\tau$

Special linear combinations are acted on by  $\hat{K}_p$  [CB, Ferrero, Zhou; 2101.04114].

$$\mathcal{M}_{\epsilon p, 0}^{(d-2)} = \mathcal{M}_{\epsilon p, 0}^{(d)} + c_{2,0}^{(d)} \mathcal{M}_{\epsilon p+2, 0}^{(d)}$$

$$\mathcal{M}_{\epsilon p, 0}^{(d-4)} = \mathcal{M}_{\epsilon p, 0}^{(d-2)} + c_{2,0}^{(d-2)} \mathcal{M}_{\epsilon p+2, 0}^{(d-2)}$$

$$= \left[ \mathcal{M}_{\epsilon p, 0}^{(d)} + c_{2,0}^{(d)} \mathcal{M}_{\epsilon p+2, 0}^{(d)} \right] + c_{2,0}^{(d-2)} \left[ \mathcal{M}_{\epsilon p+2, 0}^{(d)} + c_{2,0}^{(d)} \mathcal{M}_{\epsilon p+4, 0}^{(d)} \right]$$


$AdS_{d+1}$  and  $S^{n-1}$  dimensionality both reduce by same amount.

$$\mathcal{S}_p(x_i, t_i) = C(k_i, p) \hat{K}_p \circ \left[ W_{\epsilon p, 0}^{(d-\#Q_s/4)}(x_i) Y_p^{(n-\#Q_s/4)}(t_i) \right]$$




# More relations

Double copy is manifest for auxiliary  $AdS_5$  correlators [Zhou; 2106.07651].




# More relations

Double copy is manifest for auxiliary  $AdS_5$  correlators [Zhou; 2106.07651].




# More relations

Double copy is manifest for auxiliary  $AdS_5$  correlators [Zhou; 2106.07651].



# More relations

Double copy is manifest for auxiliary  $AdS_5$  correlators [Zhou; 2106.07651].



- Explicit Witten diagrams are being replaced by more elegant bootstrap methods.
- Some of the structures revealed by them still have a mysterious origin.
- The chiral algebra and AdS unitarity method both enable a systematic exploration of loops.
- Possible to consider both gravitons  $O(1/c_T)$  and gluons  $O(1/c_J)$  to study backreaction on the brane.
- Future targets include S-fold theories and backgrounds with defects.